

Letters

Racial and Sex Differences in Stroke Risk in Patients With Atrial Fibrillation



Atrial fibrillation (AF) is an important risk factor for ischemic stroke, but individual risk varies. Current guidelines state that AF patients without significant comorbidities are at low risk of stroke, and therefore, the benefits of anticoagulant therapy may be outweighed by the risk of bleeding (1). Although periodic reassessment of an individual's stroke risk is recommended, optimal reassessment frequency is an unsettled issue. Data from the Taiwan National Health Insurance Research Database indicate that some patients rapidly accumulate new stroke risk factors (2). However, it is unknown whether these findings translate to the United States, where AF prevalence is higher, and the population is more heterogeneous.

We sought to assess the incidence of CHA₂DS₂-VASc (congestive heart failure, hypertension, age \geq 75 years, diabetes mellitus, prior cerebrovascular accident, transient ischemic attack, or thromboembolism, vascular disease, age 65–74 years, sex) score increases in low-risk AF patients and whether score increases vary by sex or race/ethnicity. Using a validated algorithm (3), we identified 5,626 patients (mean age 66.3 ± 15.0 years) from the Mount Sinai Hospital electronic health records with newly diagnosed AF between January 2003 and December 2017 who had a CHA₂DS₂-VASc score of 0 (men) or 1 (women) at the time of AF diagnosis. This cohort was 58.5% male, 41.5% female; 51.2% Caucasian, 26.8% African American, 17.1% Hispanic/Latino, and 4.9% Asian. We computed CHA₂DS₂-VASc scores for each patient visit after AF diagnosis using a validated algorithm (4).


Cumulative incidence curves for an increase in CHA₂DS₂-VASc score from 0 to 1 (men) or 1 to 2 (women) are displayed in **Figure 1**. Overall, the CHA₂DS₂-VASc score increased in 27.4% of the study population within 1 year. At median follow-up times (2.01 years, interquartile range: 0.23 to 5.73 years), the CHA₂DS₂-VASc score increase percentages were: median: 36.0%; Q1: 15.9%; and Q3: 48.5%. Score

increases at 1 year/median (interquartile range) years occurred in 33.6/43.6 (17.1% to 59.0%) of African Americans and 30.5/38.7 (14.7% to 53.5%) of Hispanic/Latinos. Score increases occurred most frequently from new diagnosis of hypertension (62%), vascular disease (24%), and age increase (14%). A multivariable Cox proportional hazards regression model demonstrated that African Americans (hazard ratio [HR]: 1.44; 95% confidence interval [CI]: 1.33 to 1.57; $p < 0.001$), Hispanic/Latinos (HR: 1.17; 95% CI: 1.06 to 1.30; $p = 0.002$), and Asians (HR: 1.21; 95% CI: 1.02 to 1.43; $p = 0.035$) were at significantly higher risk for a CHA₂DS₂-VASc score increase relative to Caucasians. Additionally, men were at significantly increased risk relative to women (HR: 1.20; 95% CI: 1.12 to 1.29; $p < 0.001$).

Because a Class 1A recommendation exists for initiation of oral anticoagulants at CHA₂DS₂-VASc scores of 2 (men) or 3 (women) (1), we conducted a separate analysis for score increases from ≤ 1 to 2 (men) and ≤ 2 to 3 (women). In this analysis, 24.5% of the population became indicated for anticoagulation within 1 year (34.4% at 2 years). Stratification by sex and race/ethnicity revealed similar incidence ratios to **Figure 1**. African Americans (HR: 1.40; 95% CI: 1.32 to 1.49; $p < 0.001$), Hispanic/Latinos (HR: 1.29; 95% CI: 1.20 to 1.39; $p < 0.001$), and Asians (HR: 1.42; 95% CI: 1.15 to 1.60; $p < 0.001$) were at significantly higher risk for a CHA₂DS₂-VASc score increase relative to Caucasians, and men (HR: 1.23; 95% CI: 1.17 to 1.30; $p < 0.001$) were at significantly increased risk relative to women.

Although clinical guidelines recommend periodic stroke risk reassessment for AF patients, the optimal interval between reassessments is unclear. This is of particular importance in AF patients with low CHA₂DS₂-VASc scores, because risk factor acquisition may prompt oral anticoagulation therapy initiation. In our cohort of AF patients with CHA₂DS₂-VASc scores of 0 (men) or 1 (women), 27.4% acquired at least 1 new stroke risk factor within 1 year, a rate higher than previously reported (2). Furthermore, sex and race/ethnicity significantly affected stroke risk factor acquisition. Our findings align with previous reports (2) asserting stroke risk in AF increases at an alarming rate; we extend these by demonstrating differences based on sex and race/ethnicity. Our

FIGURE 1 Cumulative Incidence of CHA₂DS₂-VASC Increases From 0 to 1 (Men) or 1 to 2 (Women)



(A) Men exhibited increased risk relative to women (B); African Americans, Hispanic/Latinos, and Asians exhibited increased risk relative to Caucasians.

findings affirm the need for frequent stroke risk reassessment in AF patients, particularly for men and for patients of African American or Hispanic ethnicity. Clinicians should consider these factors when determining optimal risk reassessment intervals for their AF patients.

Adam J. Russak, MD
 Kipp W. Johnson, BS
 Jonathan L. Halperin, MD
 Bethany Percha, PhD
 *Joel T. Dudley, PhD
 *Icahn School of Medicine at Mount Sinai
 770 Lexington Avenue, 15th Floor
 New York, New York 10029
 E-mail: joel.dudley@mssm.edu
 Twitter: @jdudley, @adam_russak, @kipwjohnson, @blpercha
<https://doi.org/10.1016/j.jacc.2019.10.018>

© 2019 Published by Elsevier on behalf of the American College of Cardiology Foundation

Please note: Dr. Halperin has received consulting fees from Boehringer Ingelheim, Bayer Healthcare, Ortho-McNeil-Janssen Pharmaceuticals, and Medtronic. All other authors have reported that they have no relationships relevant to the contents of this paper to disclose. Joaquin E. Cigarroa, M.D. served as Guest Associate Editor for this paper.

REFERENCES

1. January CT, Wann LS, Calkins H, et al. 2019 AHA/ACC/HRS focused update of the 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *J Am Coll Cardiol* 2019;74:104-32.
2. Chao TF, Chiang CE, Chen TJ, Lip GYH, Chen SA. Reassessment of risk for stroke during follow-up of patients with atrial fibrillation. *Ann Intern Med* 2019;170:663-4.
3. Ritchie MD, Denny JC, Crawford DC, et al. Robust replication of genotype-phenotype associations across multiple diseases in an electronic medical record. *Am J Hum Genet* 2010;86:560-72.
4. Navar-Boggan AM, Rymer JA, Piccini JP, et al. Accuracy and validation of an automated electronic algorithm to identify patients with atrial fibrillation at risk for stroke. *Am Heart J* 2015;169:39-44.e2.